



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

from a more abundant collection. The axis was rhizomatous, giving off numerous roots at intervals, and bearing two kinds of leaves, one set of petioles being supplied by a monarch leaf trace, and the other set by a diarch trace. The smaller leaves, supplied by the monarch trace, show at base a sheathing organ which is thought to represent the so-called aphlebia of *Zygopteris*; if so, this is the first record of the structure in *Botryopteris*, and further emphasizes the relationship of the two genera. BERTRAND'S view that the simple stele of *B. antiqua* is due to reduction and not to its primitive character is objected to. As the author says, "this view involves the assumption that the diarch type of petiole is older than the monarch, and the species (*B. antiqua*) is in process of simplification. This result is not easy to harmonize with the fact that later forms of *Botryopteris* petiole are triarch."—J. M. C.

Origin of transfusion tissue.—The so-called transfusion tissue of the leaves of gymnosperms has been recognized for many years as an anatomical feature of the group. WORSDELL (1897) suggested, on the basis of distribution and nature, that it is a modified centripetal xylem. Since the presence of centripetal xylem is an important fact in discussing evolutionary sequences, this view extended the range of recognizable centripetal xylem. Now Miss CARTER³² has studied the beginnings of this tissue in the cotyledons, using 13 species, representing 9 genera of conifers. The conclusion is "that the first-formed transfusion tracheids appeared in such positions and were of such size as to make it appear improbable that they arose, in these organs at any rate, as an extension of the development of the centripetal wood." The evidence from a comparison with the other elements of the vascular strand suggests that "transfusion tissue" develops from the parenchyma.—J. M. C.

The causes of thorn development.—Since LOTHELIER conducted his researches on the experimental morphology of thorns, it has been generally believed that their development is favored and even caused by abundant light or by atmospheric desiccation. This was supposed to be the case particularly in the gorse, *Ulex europaeus*. ZEIDLER now calls these results in question,³³ for he is able to secure the development of thorns in *Ulex* both in partial darkness and in moist atmosphere. He regards the leafy shoots secured by LOTHELIER in moist air and in darkness merely as juvenile forms, whereas the thorny shoots are regarded as adult forms. It may be remarked that, even if further experiment should confirm the views of ZEIDLER, the real problem is in no wise touched by his experiments. It would still remain to determine why "juvenile shoots" should appear at some times and "adult shoots" at other times.—H. C. COWLES.

³² CARTER, M. GERALDINE, A reconsideration of the origin of "transfusion tissue." Ann. Botany 25:975-982. figs. 4. 1911.

³³ ZEIDLER, J., Ueber den Einfluss der Luftfeuchtigkeit und des Lichtes auf die Ausbildung der Dornen von *Ulex europaeus* L. Flora 102:87-95. 1911.